Agricultural water supply system resilience

Hasna MOUMNI

Karima SEBARI

Ali HAMMANI

Laila STOUR

Department of rural engineering, Hassan II Institute of Agronomy and Veterinary Medicine, Rabat, Morocco Department of rural engineering, Hassan II Institute of Agronomy and Veterinary Medicine, Rabat, Morocco Department of rural engineering, Hassan II Institute of Agronomy and Veterinary Medicine, Rabat, Morocco Laboratory of Process and Environment Engineering, Faculty of Sciences and Techniques of Mohammedia, Hassan II University of Casablanca, Mohammedia, Morocco

Resilience is an essential key of the assessment of water resources system management. Adequate description of the resilience of water resource systems needs to consider the emergent properties arising from the interaction of the component subsystems. This paper presents an approach to assess resilience in an agricultural water supply system. The system is contextualized as a meta-system composed of three subsystems; the natural catchment and reservoirs, the water distribution infrastructure and agricultural users. The proposed approach allows studying each sub-system separately to determine its properties, quantify the interdependencies between the subsystems and integrate pressures that affect the operation of each sub-system and, consequently, the system in its entirety. This work is a first step in assessing agricultural water resources' resilience under climatic and anthropogenic pressures in agricultural water resources system.

Keywords: Resilience, agricultural water resources system, climatic and anthropogenic pressures

Introduction

Water is essential for agricultural production and food security. Nevertheless, several regions in the world are facing water scarcity. In order to cover food demands of a rapidly expanding world population and improve livelihoods in rural areas, ensuring access to water is crucial (Allouche, 2011; Kang et al., 2017; McNeill et al., 2017).

Climate change will significantly impact agriculture in irrigated and rainfed areas across the globe (Lamboll et al., 2017; Shankar and Shikha, 2018). Increasing demand for water, limiting crop productivity and reducing water availability are the consequences of climate change (Ayanlade et al., 2018; Faramarzi et al., 2013). Population growth will increase water demand. Besides, urbanization and intensive agriculture are major sources of pollution of water resources. An indepth understanding of the impact of climate change and population growth on the resilience of the water supply is essential to support sustainable decision-making (McNeill et al., 2017).

Africa is particularly vulnerable to climate change (Adenle et al., 2017). This vulnerability is related to the importance of the agricultural sector for the population's livelihoods and food security and its place in the national economy (Busby et al., 2014; Filho et al., 2018). The consequences of climate change combined with the vulnerability of agriculture and livelihoods in Africa require effective climate change mitigation and adaptation measures in the region (Epule et al., 2017; Nguimalet et al., 2016). The Paris Agreement recognizes that adapting agriculture to climate change is closely linked to water management. Adaptation measures will enhance both resilience and water security (Tobin et al., 2018). To capitalize on the Paris agreement's efforts, the Cop 22 held in Morocco aimed to unblock the financing of climate change adaptation measures by focusing mainly on the

developing countries most vulnerable to climate change. Launched on this occasion, the Triple A initiative: "Adaptation of African Agriculture" aims to reduce Africa and agriculture's vulnerability to climate change (Ghezloun et al., 2017).

Reservoirs play a crucial role in securing water in regions suffering from water scarcity. Many regions rely entirely on reservoirs for water supply, particularly where groundwater resources are limited, climate variability is high and alternative sources of water are not available (Mereu et al., 2016). Climate change and increasing water demand are major challenges in these reservoirs' management and resilience (Zhang et al., 2017; Zhao et al., 2018). A better understanding of the impacts of these pressures on the water balance of the reservoirs is necessary.

An agricultural water supply system is complex. It consists of a supply catchment and storage reservoir, a distribution system and agricultural users, ranging from socio-ecological to technical domains. Due to the complexity of the agricultural water system and the multiple interactions among its components, different management approaches need to be applied to ensure a reliable supply. The system needs to be highly resilient to withstand the pressures of climate change and population growth.

Assessment of water supply system performance needs indicators that provide pertinent information on system performance. Resilience is increasingly recognized as a key system property of underpinning sustainability (Blackmore and Plant, 2008) but transforming the concept of resilience into an operational tool for policy and management purposes is a challenge (Klein et al., 2003).

Resilience as a management concept was explored by Amarasinghe et al., (2012) and Barnes et al., (2012) for application in the potable water supply in Southeast Queensland, an area with one of Australia's most significant urban growth rates. They explored an approach to modeling surrogates of resilience as related families of disaggregated models and considering the viability of these models as decision support aids in managing the complex interaction between components inherent in a potable water supply system.

This paper seeks to examine the hypothesis that resilience is an essential management strategy in agricultural water supply management. The approach proposed here, adapted from Amarasinghe et al., (2012) and Barnes et al., (2012), aims to study a system of agricultural water supply as a system of systems. The work aims to identify dependencies and interconnection between system components and then disaggregate these relationships to define parameters that describe the behavior of the system. These parameters can be developed as indicators describing the resilience.

The Moroccan context and the water resources system of Ouergha

Morocco, like many other developing countries, is strongly impacted by climate change. Water resources are characterized by scarcity and irregularity. They are subject to increasing pressure linked to population growth, to development of irrigated agriculture as well as to urban, industrial and tourism development. Climate change impacts and increasing water demand, would make meeting the water needs more complex (Schilling et al., 2012; Seif-Ennasr et al., 2016). Studies of the evolution of climatic factors have demonstrated an increase in temperatures and an accentuation of climatic variability, mainly characterized by a change in the intensity and distribution of rainfall, a reduction in water resources and an amplification of extreme events (El Jihad et al., 2014; Khattabi et al., 2014; Stour and Agoumi, 2008; Tramblay et al., 2012).

The Moroccan economy will suffer from this situation because it is closely dependent on the abundance of rainfall, as highlighted by the strong dependence between GDP and agricultural GDP. Climate change is affecting the economic development of many sectors, mainly agriculture.

Given these constraints, the development of water resources has always been a central concern of the country's economic policies, particularly because of their strategic role in developing irrigated agriculture and the country's water and food security. Since the beginning of the 1960s, Morocco had adopted a strategy that provided the country with extensive hydraulic infrastructure and ensured water needs (drinking water, flood protection, and agricultural production).

The Ouergha basin (7200 Km2 upstream of the M'jaara hydrometric station) regulates almost a third of Morocco's surface water resources (Senoussi et al., 1999). It feeds the Al Wahda dam, the largest reservoir in the country and the second-largest in Africa in terms of storage capacity with 3.7 billion m3. Rainfall can reach 1800 mm/year in mountainous areas; the country's wettest areas. The bedrock of the basin is impermeable (clay and marl) preventing infiltration and development of groundwater resources (Boukrim et al., 2011; Senoussi et al., 1999).

Boukrim's work on the impact study of climate change showed an annual warming of 0.15oC and a rainfall deficit of about 4 mm observed from 1982/1983 to 2007/2008 which is the basin's driest period (Boukrim et al., 2011). As a result of the increase in temperature, more evaporation reduced flow into rivers and the active storage volume of reservoirs. Senoussi et al., (1999) assessed the potential impacts of global warming in the case of a doubling of the atmospheric CO2 concentration on freshwater potential at the Al Wahda Dam. The results obtained indicate a decrease in annual runoff at this basin reaching 8% by 2050. This is a significant deficit in a country where the resource is particularly scarce.

The influence of climatic and anthropogenic pressure on water resources in a basin where agricultural activities continue to increase and feeds important agricultural areas (such as Sais and the Gharb plain) must be considered. In the absence of groundwater resources, it is important to improve surface water resources management tools as it is the sole water source of this basin. As resilience is a key to good management, this work is a first step in assessing agricultural water resources' resilience under climatic and anthropogenic pressures in this water resource system.

Resilience of water supply systems

Measuring the resilience of a WRS is an important consideration in the management of water resources system (Li and Lence, 2007). Traditional water management approaches have failed to achieve sustainable results as they have treated social systems and ecosystems as distinct systems (Bohensky, 2006). WRS cannot be sustained by maintaining the individual components making up the system (Liu et al., 2012). WRS as social-ecological systems acts as a strongly coupled, complex and evolving integrated systems (Cofie and Amede, 2015).

Resilience is a measure of how fast a system is likely to return to a satisfactory state once the system has entered an unsatisfactory state (Kjeldsen and Rosbjerg, 2004). There have been several studies on developing quantitative metrics of water resource systems resilience since the 1960s. The most widely used and cited definition may be the one by Hashimoto et al., (1982). They defined resilience as the average probability of recovery at time step t+1 from a failure state at time step t. In this sense, resilience represents the rapidity of the system returning to a satisfactory state after an occurrence of failure. Moy et al., (1986) used the maximum time duration of failure as a measure of resilience; the longer the duration, the less resilient the system is. In this sense, resilience is defined as the maximum duration that the system stays in an unsatisfactory state.

Other definitions of water supply systems' resilience are found in the literature (Kjeldsen and Rosbjerg, 2004; Wang and Blackmore, 2009). Most of these definitions are focused on quantifying general resilience. The system's specific resilience to a particular type of pressures such as climate change or population growth was not considered. Specified resilience – resilience "of what to what" is more concrete and amenable to assessment in contrast to general resilience (Carpenter et al., 2001). Most of these definitions used one parameter for quantifying resilience. As a Meta system,

considering only one parameter to assess the resilience is difficult (Amarasinghe et al., 2012). These definitions may be helpful in the context of traditional reservoir management but not for a complex system of subsystems. The development of an effective resilience assessment methodology is a major challenge because of the complexity of the water supply system and the different types of pressures that can affect the system (Haimes, 2012).

Modeling agricultural water supply system

As a system of systems

The first step in defining systemic resilience is a in-depth understanding of the system and the interconnections between its subsystems. The approach taken here is to define a complete agricultural water supply system. A complete agricultural water supply system consists of water catchment and the reservoirs, water distribution infrastructure and agricultural users. Integration of the three components forms a system of systems as described in Figure 1. The Metasystem is complex since its subsystems belong to different domains ranging from sociological, technical to the ecological domain. Assessment of the resilience of the agricultural water supply system requires consideration of the three components.

Climate change and population growth are two major sources of pressure on the agricultural water supply system. They have a significant impact on the quality and quantity of water in the reservoir storage.

The resilience of agricultural water supply system

The approach presented here is exploratory. It is based on a in-depth conceptual understanding of the agricultural water supply infrastructure. In developing the first expression, the system's three components have been considered and examined from upstream to downstream.

First-degree expression

Rs = f(QT, QD, DM, a)(1)

Rs: Resilience of supply,

QT: Water quantity,

QD: Distributed water,

DM: Agricultural water demand,

a: Other variables.

Second-degree expression

QT = f (
$$\Sigma$$
 [I - O], Σ L, b) (2)

I: All inflows into the system 1 (Surface runoff, groundwater flow, diversions, transfers and others. In Ouergha context, it consists mainly of surface runoff),

O: All outflows from the system 1 (Domestic demand, environmental flows and others),

 Σ L: Summation of all losses (Percolation, evapotranspiration, leakage or other accounted losses and reservoir overflows during high rainfall),

b: Other variables.

 $QD = f(\Sigma L, c)(3)$

 Σ L: Summation of all losses in system 2 (Evaporation, leakage and overflow of the irrigation channels),

c: Other variables.

Dm = f(Bni, Si, Eq, d) (4)

Bni: Net irrigation water requirements in system 3,

Si: Area occupied by crops,

Eg: Overall irrigation efficiency,

d: Other variables.

The initial conceptual model is decomposed into a series of cascading empirical models of reducing complexity. The proposed approach will allow studying each subsystem separately to determine its properties, quantify the interdependencies between the subsystems, integrate the climatic and anthropic pressures that affect the operation of each sub-system, and consequently the system in its entirety.

Conclusion

This work presents an approach to modeling the resilience of an agricultural water supply system. The system consists of water catchment and the reservoir, water distribution infrastructure and agricultural users. The water system was considered as system of systems to appreciate the complexity of Agricultural WSS. This work is a first step for developing a suite of indicators and metrics to define resilience of an agricultural water supply system to climate change and population growth. The proposed modeling approach needs to be further developed to improve agricultural water supply systems management in a resilient context.

References

Adenle A.A., Ford J.D., Morton J., Twomlow S., Alverson K., Cattaneo A., Cervigni R., Kurukulasuriya P., Huq S., Helfgott A., Ebinger J.O. (2017). Managing climate change risks in Africa - A global perspective. Ecol. Econ., 141: 190-201.

Allouche J. (2011). The sustainability and resilience of global water and food systems: Political analysis of the interplay between security, resource scarcity, political systems and global trade. Food Policy, The challenge of global food sustainability, 36: S3-S8.

Amarasinghe P., Barnes P., Egodawatta P., Goonetilleke A. (2012). Application of Resilience concept for enhanced management of water supply systems, in: Proceedings of the 2nd International Conference on Sustainable Built Environment, Kandy, Sri Lanka.

Ayanlade A., Radeny M., Morton J.F., Muchaba T. (2018). Rainfall variability and drought characteristics in two agro-climatic zones: An assessment of climate change challenges in Africa. Sci. Total Environ., 630: 728-737.

Barnes P., Egodawatta P., Goonetilleke A. (2012). Modelling resilience in a water supply system

contrasting conditions of drought and flood, in: Kakimoto, R., Yamada, F. (Eds.), In: International Conference on Disaster Management(IIIRR 2012), Kumamoto University, Kumamoto, Japan, pp. 339–346.

Blackmore J., Plant R. (2008). Risk and resilience to enhance sustainability with application to urban water systems. J. Water Resour. Plan. Manag.-Asce, 134.

Bohensky E.L. (2006). A Social-Ecological systems perspective on water management in South Africa (Thesis). University of Pretoria.

Boukrim S., Sadkaoui N., Lahrach A., Chaouni A. (2011). Etude d'impact des changements climatiques sur les ressources hydriques du bassin versant de l'Ouergha (Rif-Maroc). J. Eau Environ., 10: 42–55.

Busby J.W., Smith T.G., Krishnan N. (2014). Climate security vulnerability in Africa mapping 3.01. Polit. Geogr., Special Issue: Climate Change and Conflict, 43: 51-67.

Carpenter S., Walker B., Anderies J., Abel N. (2001). From metaphor to measurement: Resilience of what to what? Ecosystems, 4: 765-781.

Cofie O., Amede T. (2015). Water management for sustainable agricultural intensification and smallholder resilience in sub-Saharan Africa. Water Resour. Rural Dev., Managing Rainwater and Small Reservoirs in Sub-Saharan Africa, 6: 3–11.

El Jihad M.-D., Peyrusaubes D., El Bouzidi A. (2014). Seasonal droughts and climate change in the Gharb (Morocco). Ruralités, Le Gharb, un territoire à l'épreuve du changement climatique. Coordination: M-D. El Jihad et D. Peyrusaubes (Eds.), 14-25.

Epule, T.E., Ford, J.D., Lwasa, S., Lepage, L. (2017). Climate change adaptation in the Sahel. Environ. Sci. Policy 75, 121-137.

Faramarzi M., Abbaspour K.C., Ashraf Vaghefi S., Farzaneh M.R., Zehnder A.J.B., Srinivasan R., Yang H. (2013). Modeling impacts of climate change on freshwater availability in Africa. J. Hydrol., 480: 85–101.

Filho W.L., Balogun A.-L., Ayal D.Y., Bethurem E.M., Murambadoro M., Mambo J., Taddese H., Tefera G.W., Nagy G.J., Fudjumdjum H., Mugabe P. (2018). Strengthening climate change adaptation capacity in Africa- case studies from six major African cities and policy implications. Environ. Sci. Policy, 86: 29–37.

Ghezloun A., Saidane A., Merabet H. (2017). The COP 22 New commitments in support of the Paris Agreement. Energy Procedia, International Conference on Technologies and Materials for Renewable Energy, Environment and Sustainability, TMREES17, 21-24 April 2017, Beirut Lebanon 119: 10-16.

Haimes Y.Y. (2012). Modeling complex systems of systems with Phantom System Models. Syst. Eng., 15: 333–346.

Hashimoto T., Stedinger J., Loucks, P. (1982). Reliability, resiliency, and vulnerability criteria for water resource system performance evaluation. Water Resour. Res., 18: 14-20.

Kang S., Hao X., Du T., Tong L., Su X., Lu H., Li X., Huo Z., Li S., Ding R. (2017). Improving agricultural water productivity to ensure food security in China under changing environment: From research to practice. Agric. Water Manag., 179: 5–17.

Khattabi A., Chriyaa A., Hammani A., Moudoud B. (2014). Vulnérabilités climatiques et stratégies de développement: Synthèse et recommandations stratégiques pour une prise en compte du risque «climat» dans les politiques et stratégies sectorielles. Technical report . Institut Royal des Etudes Stratégiques (IRES), Rabat.

Kjeldsen T.R., Rosbjerg D. (2004). Choice of reliability, resilience and vulnerability estimators for risk assessments of water resources systems. Hydrol. Sci. J., 49: 767.

Klein R.J.T., Nicholls R.J., Thomalla, F. (2003). Resilience to natural hazards: How useful is this concept? Glob. Environ. Change Part B Environ. Hazards, 5: 35–45.

Lamboll R., Stathers T., Morton J. (2017). Chapter 13 - Climate Change and Agricultural Systems, in: Snapp, S., Pound, B. (Eds.), Agricultural Systems (Second Edition). Academic Press, San Diego, pp. 441–490.

Li Y., Lence B.J. (2007). Estimating resilience for water resources systems. Water Resour. Res. 43(7).

Liu D., Chen X., Nakato T. (2012). Resilience Assessment of Water Resources System. Water Resour. Manag., 26: 3743–3755.

McNeill K., Macdonald K., Singh A., Binns A.D. (2017). Food and water security: Analysis of integrated modeling platforms. Agric. Water Manag., 194: 100–112.

Mereu S., Sušnik J., Trabucco A., Daccache A., Vamvakeridou-Lyroudia L., Renoldi S., Virdis A., Savić D., Assimacopoulos D. (2016). Operational resilience of reservoirs to climate change, agricultural demand, and tourism: A case study from Sardinia. Sci. Total Environ., 543: 1028–1038.

Moy W.-S., Cohon J.L., ReVelle C.S. (1986). A Programming model for analysis of the reliability, resilience, and vulnerability of a water supply reservoir. Water Resour. Res., 22: 489-498.

Nguimalet C.-R., Mahe G., Laraque A., Orange D., Yakoubou M. (2016). Note sur le changement climatique et gestion des ressources en eau en Afrique: repenser l'usage et l'amélioration des services éco-systémiques de l'eau. Geo-Eco-Trop 40: 317-326.

Schilling J., Freier K.P., Hertig E., Scheffran J. (2012). Climate change, vulnerability and adaptation in North Africa with focus on Morocco. Agric. Ecosyst. Environ., 156: 12–26.

Seif-Ennasr M., Zaaboul R., Hirich A., Caroletti G.N., Bouchaou L., El Morjani Z.E.A., Beraaouz E.H., McDonnell R.A., Choukr-Allah R. (2016). Climate change and adaptive water management measures in Chtouka Aït Baha region (Morocco). Sci. Total Environ., 573: 862–875.

Senoussi S., Agoumi A., Yacoubi M., Fakhraddine A., Sayouty E.H., Mokssit A., Chikri N. (1999). Changements climatiques et ressources en eau Bassin versant de l'Ouergha (Maroc). Hydroécologie Appliquée, 11: 163–182.

Shankar S., Shikha (2018). Chapter 7 - Impacts of Climate Change on Agriculture and Food Security, in: Singh, R.L., Mondal, S. (Eds.), Biotechnology for Sustainable Agriculture. Woodhead Publishing, pp. 207–234.

Stour L., Agoumi A. (2008). Sécheresse climatique au Maroc durant les dernières décennies. Hydroécologie Appliquée, 16: 215–232.

Tobin P., Schmidt N.M., Tosun J., Burns C. (2018). Mapping states' Paris climate pledges: Analysing targets and groups at COP 21. Glob. Environ. Change, 48: 11-21.

Tramblay Y., Badi W., Driouech F., El Adlouni S., Neppel L., Servat E. (2012). Climate change impacts on extreme precipitation in Morocco. Glob. Planet. Change, 82–83: 104–114.

Wang C.-H., Blackmore J. (2009). Resilience Concepts for Water Resource Systems. J. Water Resour. Plan. Manag.-Asce - 135.

Zhang W., Liu P., Wang H., Lei X., Feng M. (2017). Operating rules of irrigation reservoir under climate change and its application for the Dongwushi Reservoir in China. J. Hydro-Environ. Res., 16: 34-44.

Zhao G., Gao H., Kao S.-C., Voisin N., Naz B.S. (2018). A modeling framework for evaluating the drought resilience of a surface water supply system under non-stationarity. J. Hydrol., 563: 22–32.

References